

Jazzy

Revolutionising the salad potato market

Jazzy is a second early variety that produces very high numbers of white skinned long oval tubers with light yellow flesh, well suited to the production of a quality punnet crop. Jazzy tubers have excellent culinary qualities, can be cooked in a variety of different ways, and can be utilised from fresh lifted, and throughout the storage season. Resistance to Common Scab is good, although Jazzy is susceptible to Foliage Blight and PCN.

	Jazzy
Yield	High
Underwater Weight	340
Dry Matter %	18.8
Maturity	Early/Medium
Dormancy	Very Short
Bruising index (*1)	6
Second Growth (*2)	6.7
Common scab (*2)	6.6
Late Blight (*2)	2.8
Tuber Shape	Long
Percentage 50/+	19
Percentage 40/70	61
Skin Colour	Light Yellow
Flesh Colour	Light Yellow
Eye Depth	Very Shallow
Uniformity	Good
Washability	Very Good
Internal Defects (%)	1
Hollow Heart (%)	0
Cooking type February	Firm
Fry Colour fries October (8 °C)	4
Fry Colour fries February (8 °C)	3
Fry Colour fries May (8 °C)	3
Potato Cyst Nematodes Ro1 (*2)	3
Potato Cyst Nematodes Ro2/3 (*2)	
Potato Cyst Nematodes Pa2 (*2)	
Potato Cyst Nematodes Pa3 (*2)	
Wart Disease wz 1 (D1) (*3)	10
Wart Disease wz 2 (G1) (*3)	7
Wart Disease wz 6 (O1) (*3)	7
Wart Disease wz 18 (T1) (*3)	8

(*1): 0= none, 50 = severe

(*2): 9 = resistant, 1 = very susceptible

(*3): 10 = resistant, 3 = very susceptible

Characteristics

- Extremely numerous and even sized
- Excellent culinary properties
- Low risk of bruising
- Can be stored throughout the season



Agronomy

Soil Nitrogen Supply*	very low, low	medium	high, very high
N input kg/ha	100-160	60-120	40-80

Soil Supply*	very low	low	medium	high	very high
P input kg/ha	250	210	170	100	0
K input kg/h	360	330	300	150	0
Mg input kg/ha	120	80	40	0	0

Storage	1.5°C/3°C	8/6 months
---------	-----------	------------

50 kg Tuber Count	500	700	900	1100	1300	1500	1700
Population/ha	63773	79101	89716	98776	108638	118540	125008
Spacing/cm	28,85	20,84	18,38	16,69	15,17	13,91	13,19
Seed Rate t/ha	6,38	5,65	4,99	4,49	4,18	3,96	3,68

Values are based on an average soil type and planting date.

We recommend to contact your local agronomist and adjusting to your own situation and experience.

Timing	Pre-em	Post-em
Metribuzin sensitivity	None	Slightly

General advice for best results

- Best cultivated in bed system if possible to maximise homogenous tuber sizing
- Ensure a robust blight programme
- Store relatively cold as Jazy has a very low dormancy

For any further information
contact your local Meijer
representative or visit our
website at:
www.agriana.pl

AGRIANA 

Agriana Sp. z o.o.
ul. Główna 9
82-224 Szymankowo
NIP: 579-22-53-712
Tel.: +48 55 271 19 84

Every effort has been made to ensure that the advice, information, recommendations and statements given in this management guide are correct. Agriana Sp. z o.o. and associated companies cannot be held responsible for any damage or loss resulting from their use.