








Added value of the variety

Foliar late blight resistant. 
Tasty table potato with low input requirement.
Washable and attractive skin.
Gives high yields from all soil types.
Tolerant to climatic stress.













Characteristics


Shape of tuber	RO	
Av. Tuber length (Ø 50-60 mm)	7,5 cm	
Maturity (9=early, 4=late)	6,6	
Colour of skin	LY	
Colour of flesh	6,2	
Suitability for cooking	7,8 A	

Soil adaptability

	Sandy	Light	Heavy
Yield (Relative)	103	99	100
DM (%)	17,3	18,2	19,0
UWW	311	330	347
Tuber size (Ø mm)	55	59	62

Ro1	9	
Ro2/3	6	
Ro5	2	
Pa2	1	
Pa3	1	
wz 1 (D1)	10	
wz 2 (G1)	4	
wz 6 (O1)	4	
wz 18 (T1)	6	
TRV	5	
Moptop virus	Rather susceptible	




Climate adaptability

Drought tolerance	Tolerant
Secondary growth	7 






Fertiliser requirement

	N	P	K	Mg	B	Ca	Mn
Quantity	Low (-25%)	Standard	Standard	Standard	Standard	Standard	Standard
No. applications	1	1	1	1	1	1	1

Chemical requirement







Foliar late blight	9 
Tuber late blight	Rather tolerant
Early blight	Rather tolerant
Y-virus	8 
Yntn-virus	10 
Leafroll virus	Rather susceptible
Black dot	Rather susceptible
Silver scurf	Rather susceptible
Metribuzin pre-emergence	Susceptible
Metribuzin post-emergence	Tolerant

Storability

Dormancy (2=short; 8=long)	4,9 
Pressure spots	Rather susceptible
Fusarium	4 
Frying /crisping colour in October*	5 
Frying /crisping colour in February*	4 
Frying /crisping colour in May*	5 

*Potatoes stored at 8 °C

Marketable yield

Bruising index (0-50)	11 
Washability	7 
Common scab	6 
Powdery scab	7 
IRS	2,7 
Hollow heart	0,7 
Mechanical Damage	Rather tolerant



Planting


Before planting keep the seed quiet; plant them out of cold storage.
Do not cultivate on soils prone to common scab.
Plant deep and make big ridges in order to avoid tuber greening.
Plant in warm soils in order to avoid rhizoctonia.

Size	75 cm	90 cm	Plant/ha
28/30	22 cm	18 cm	60600
30/40	28 cm	23 cm	47600
40/50	36 cm	30 cm	37000
50/55	37 cm	31 cm	36000



Fertiliser input (kg/ha):

Depending on the soil supply and applicable national legislation; total amount required by the variety.



Nitrogen:	(N)	180	180
Phosphate:	(P ₂ O ₅)	110	110
Potassium:	(K ₂ O)	250	250

To prevent mineral deficiencies, apply these elements in time (B, Mg, Mn, S)

Supply 25% less nitrogen than your references varieties and apply all at planting time.



Crop protection

Susceptible to the use of 450 g/ha of Metribuzin at pre-emergence.
Take care of the late blight foliar resistance, do not neglect your blight program.



Harvest and storage

Wait for a proper set skin before harvest.
Short dormancy; after harvest and wound healing cool down quickly keep the temperature stable.
Start sprouting inhibitor application as soon as potatoes are dry.
Susceptible to Fusarium.



Storage:

3 °C
6 months

Key advice to succeed this variety:

Before planting keep the seed quiet; plant them out of cold storage.
Plant deep and make big ridges in order to avoid tuber greening.
Supply 25% less nitrogen than your references varieties and apply all at planting time.
Wait for a proper set skin before harvest.
Start sprouting inhibitor application as soon as potatoes are dry.

Avoid
desprouting
before
planting